


1999-2003



# Community Health Profiles



## Greater Hobby Area Super Neighborhood



*Providing Health Information  
for Community Action*

# Introduction



This community health profile highlights important health issues facing the residents of the Greater Hobby Area Super Neighborhood.

In Houston, a “super neighborhood” is a geographically defined area where residents, civic organizations, institutions and businesses work together to identify, plan, and set priorities to address the needs and concerns of their community. The boundaries of each super neighborhood rely on major physical features such as bayous or freeways to group together contiguous communities that share common physical characteristics, identity or infrastructure. Greater Hobby Area Super Neighborhood will hereinafter be referred to as “Greater Hobby Area”.

It is the intention of the Houston Department of Health and Human Services (HDHHS), in developing health profiles such as this, to promote a better understanding by local residents, community-based organizations, community leaders, medical providers, and the public health community of the unique character and circumstances of our various communities, and to draw attention to those matters that contribute to the greatest of health disparities among the citizens of our growing, culturally and ethnically diverse city.

This profile also represents an effort on the part of HDHHS to provide a “baseline” of indicators of health in our communities, against which future trends in conditions can be measured and monitored, and appropriate public health actions, taken.

We hope that this health profile will support these efforts in Greater Hobby Area and across the City of Houston.


## Table of Contents

Community Resources.....	3
The Super Neighborhood at a Glance.....	4
Major Causes of Death.....	6
Years of Potential Life Lost.....	7
Maternal and Child Health.....	8
Births to Teen Mothers.....	9
Infant Mortality.....	10
Leading Causes of Hospitalization.....	11
Crime.....	12
Tuberculosis.....	13
Drowning and Submersions.....	13
Food-related Illness.....	13
Environmental Health & Safety.....	14
HIV/AIDS.....	15
Gonorrhea.....	16
Syphilis.....	17
Chlamydia.....	18
Technical Notes.....	19

**Stephen L. Williams, M.Ed., M.P.A.**  
**Director**  
**Houston Department of Health and Human Services**

# Community Resources

The health of a community depends to a great extent upon the availability and accessibility of its resources.




# Greater Hobby Area at a Glance

The total population of Greater Hobby Area was 41,198, according to the 2000 census.\*

## Age


At the time of the 2000 census, more than one-third (35%) of Greater Hobby Area residents were under the age of 20. More than half (58%) were between 20 and 64 years of age, and 7% were 65 or older.



## Race, Ethnicity, National Origin


Approximately half of the residents in Greater Hobby Area were Hispanic. Blacks were the second largest ethnic group, comprising one-quarter of the population. Twenty-four percent of the population were of other races.

Of the total population, more than half (57%) were native Texans; 28% were foreign born.



## Employment

More than half of Greater Hobby Area residents, ages 16 and over, were employed in 1999.




\* Data Source: U.S. Census 2000. Total population was calculated from census block-level data using Summary File 1. For purposes of describing demographics using Summary File 3, the super neighborhood is defined by the following census geographies: Tracts 3331 through 3338.

### Poverty

Seventeen percent of the population in Greater Hobby Area was below the poverty level in 1999. Forty-four percent of all residents in the super neighborhood had incomes less than twice the poverty level.

Of those living below the poverty level, 43% were children under 18 years of age; 3% were adults 65 and older.





### Education

More than one-third (39%) of Greater Hobby Area residents, ages 25 and over, reported that they had not graduated from high school.

One-quarter of residents reported a high school diploma (or the equivalent) as their highest level of educational attainment.

Approximately 35% of residents had attained education beyond the high school level, with 14% earning a college degree.

\*Due to rounding, the total percentages may not be equal to 100.



### Population Stability

Nearly half of the residents of Greater Hobby Area had lived in the same house since 1995. More than one-third moved to Greater Hobby Area from other locations in Harris County between 1995 and 1999.

Fifteen percent of residents moved to the area from outside Harris County between 1995 and 1999.

Data Source: U.S. Census 2000, Summary File 3

## Major Causes of Death

During the years 1999-2003, the residents of the super neighborhood had a lower overall annual average mortality rate than that of Houston as a whole. However, the mortality rates for heart disease, cancer, and diabetes mellitus were higher in the community than Houston overall.

### Leading Causes of Mortality, Greater Hobby Area, Houston, Texas, 1999-2003

Rank	Cause of Death	Greater Hobby Area		Houston	Greater Hobby Area - Houston
		Deaths	Rates*	Rates*	Rates
	All Causes	1100	879.5	898.2	-18.7
1	Heart Disease	301	265.3	262.0	3.3
2	Cancer	260	208.0	197.6	10.4
3	Stroke	70	61.9	76.0	-14.1
4	Accidents	54	32.0	34.8	-2.8
5	Chronic Lower Respiratory Disease	34	29.6	31.9	-2.3
6	Diabetes Mellitus	34	29.3	28.0	1.3
7	Influenza and Pneumonia	24	--	20.0	--
8	Alzheimer's Disease	19	--	20.5	--
9	Septicemia	21	--	18.1	--
10	Kidney Disease	19	--	15.8	--

### Other Causes of Death of Particular Interest, Greater Hobby Area, Houston, Texas, 1999-2003

Cause of Death	Greater Hobby Area		Houston	Greater Hobby Area - Houston
	Deaths	Rates*	Rates*	Rates
Coronary Heart Disease	194	171.0	174.1	-3.1
Bronchus-Lung Cancer	68	56.0	52.8	3.2
Motor Vehicle Accident	23	--	13.2	--
Firearm Related	21	--	7.4	--
Drug-Induced Cause	13	--	8.2	--
Cervical Cancer	<5	--	2.2	--

\*Age-adjusted mortality rates: annual average deaths per 100,000 population; census 2000 populations as the denominators; age-adjusted to the 2000 US Standard Million; deaths with known age and disease information.

-- Numbers of deaths were too small for rate calculation.

Data Sources: Texas Department of State Health Services, Vital Statistics; US Census, 2000

# Years of Potential Life Lost (YPLL)

Years of Potential Life Lost (YPLL) is an indicator of premature mortality. This indicator suggests social and economic loss owing to premature death. It also gives information on the specific causes of deaths affecting younger age groups.

Leading Causes of Premature Death	YPLL Rate*	YPLL Rate**	Houston YPLL Rate**
Cancer	727.7	853.9	816.3
Accidents	693.8	640.0	779.0
Homicide	579.3	514.9	407.5
Heart Disease	413.5	551.3	689.3
Conditions Originating in the Perinatal Periods	395.2	--	-
Congenital Disorders	329.4	--	-
HIV/AIDS	239.4	--	-
Suicide	180.4	--	-
Stroke	137.5	--	-
Chronic Liver Disease-Cirrhosis	125.5	--	-
<b>Specific Causes of Interest</b>			
Firearm Related	370.1	--	-
Motor Vehicle Accident	340.9	--	-
Coronary Heart Disease	225.3	315.1	376.1
Drug-Induced Cause	167.8	--	-
Bronchus-Lung Cancer	61.7	--	-

NOTE: Special cause of death categories may not be mutually exclusive.  
 \* Crude annual average YPLL per 100,000 population under age 65 years.  
 \*\* Age-adjusted annual average YPLL per 100,000 population under age of 65, standardized for 2000 US Standard Million.  
 -- Number of deaths too small for age-adjustment.  
 - Houston data not presented because comparison data were not available for the community.

## Differences in YPLL rates between Men and Women, 1999-2003

Premature deaths from accidents, homicide, heart disease, and perinatal period conditions had higher impact on annual average YPLL rates among males than females in the community, while premature deaths from cancer had higher impact in females than males.

## Rate of Years of Potential Life Lost (YPLL Rate)

At every age of death, there is a certain number of years of "expected life" that are not lived, and are therefore "lost". The amount of lost years of life often differ by cause of death. Many people consider death before the age of 65 years as premature. More years of life were lost prematurely due to cancer, accidents, homicide, and heart disease related deaths in this community than any other causes.

The age-adjusted annual average YPLL rates for cancer and homicide were higher in the super neighborhood, while the rates for accidents and heart disease were lower compared to Houston overall. Comparison of other age-adjusted YPLL rates is not possible due to the relatively smaller number of deaths occurring before age 65 in Greater Hobby Area. YPLL is not reported where fewer than 5 deaths occurred.

Leading Causes of Premature Death §	Male YPLL Rates (number of deaths)	Female YPLL Rates (number of deaths)
Accidents	1149.8 (32)	228.2 (7)
Homicide	722.4 (21)	433.2 (10)
Cancer	659.2 (49)	797.7 (45)
Heart Disease	529.9 (44)	294.8 (23)
Conditions Originating in the Perinatal Periods	456.4 (7)	332.8 (5)
<b>Specific Causes of Interest</b>		
Motor Vehicle Accident	590.9 (17)	
Firearm Related	462.6 (14)	275.8 (7)
Coronary Heart Disease	303.2 (31)	145.8 (11)

§ Ranked by Male YPLL Rate  
 Note: Annual average YPLL rates might be unstable due to small number of premature deaths.


Data Sources: Texas Department of State Health Services, Vital Statistics; US Census, 2000

# Maternal and Child Health

Prenatal care is the care a woman gets during pregnancy. Both prenatal care and birth weight are good indicators of a newborn's chances of survival, growth, long term health, and psycho-social development.


## Entry into Prenatal Care by Trimester of Pregnancy, 1999-2003

A higher proportion of Asian women (83.3%) than Hispanic (75.7%), White (75.6%), and Black (75.0%) women in Greater Hobby Area entered prenatal care during the first trimester. A small proportion of women in all groups entered prenatal care very late in their pregnancy, or received no care at all.



## Low Birth Weight Births (LBWB), 1999-2003

Approximately 7% of live births in Greater Hobby Area had low birth weight (2500 grams or less), which was lower than Houston as a whole. Both were above the Healthy People 2010 goal of less than 5% of live births being low weight.





Low birth weight is a factor significantly related to infant mortality. Infants born with low birth weights are at increased risk for serious health problems and long term disabilities such as mental retardation, cerebral palsy, and respiratory, vision, and hearing problems. Low birth weight and infant mortality are therefore among the most important indicators of a community's health.

Data Source: Texas Department of State Health Services, Vital Statistics, 1999-2003

## Births to Teen Mothers

Teenage childbearing is associated with negative consequences for the children born of teen mothers. In addition, there are important social and economic costs to individuals as well as the society as a result of births to teenage mothers.




### Births by Age of Mother, 1999-2003

A total of 4,663 births were recorded over the period 1999-2003 among mothers in Greater Hobby Area. One out of every 7 of these births was to a young mother (10-19 years of age).


### Births to Teen Mothers, 1999-2003

The annual average birth rate for 15-17 year-old teens in Greater Hobby Area (48.1 per 1,000 females aged 15 to 17 years) was comparable to the rate in Houston overall. The birth rate among 18-19 year-old females in Greater Hobby Area was slightly higher than the total Houston rate.



### Births to Teen Mothers by Super Neighborhood, 1999-2003

Greater Hobby Area was among the neighborhoods in Houston with the lowest annual average rates of births to teen mothers (15-17 years of age).





Data Sources: Texas Department of State Health Services, Vital Statistics; US Census 2000

# Infant Mortality

Infant mortality annual average rate is the death of infants in the first year of life. It is one of the most important indicators of the health of a community. The Healthy People 2010 goal is to eliminate disparities among racial and ethnic groups with infant mortality rates (IMR) above the national average. The targeted groups are African American, American Indian, Alaskan Native and Puerto Rican populations.

## Infant Mortality Rate, 1999-2003

The annual average infant mortality rate in Greater Hobby Area was comparable to Houston's IMR, but 38% higher than the Healthy People 2010 goal of 4.5 infant deaths per 1,000 live births. Fifty-five percent (55%) of all infant deaths were among Hispanics in this community. The annual average IMR among Hispanics was slightly higher than that of Hispanics in Houston overall. However, the annual average IMR among Blacks in Greater Hobby Area was slightly lower than that of Blacks in Houston as a whole. Infant mortality rate among other races/ethnicities was not reported due to small number of infant deaths.



## Infant Mortality Rate by Super Neighborhood 1999-2003

When compared to the Healthy People 2010 goal, Greater Hobby Area was among the neighborhoods in Houston with high annual average infant mortality rates.

Data Source: Texas Department of State Health Services, Vital Statistics

# Leading Causes of Hospitalization

Much of the information on health issues that the super neighborhood residents face on a daily basis is not readily available. The leading causes of hospitalization provide a partial picture of those conditions.

Principal Diagnosis, Multiple Level Clinical Classification of ICD 9	Counts
<b>1 Complications of pregnancy; childbirth; and the puerperium</b>	<b>3489</b>
Complications mainly related to pregnancy	956
Indications for care in pregnancy; labor; and delivery	800
Complications during labor	795
<b>2 Certain conditions originating in the perinatal period</b>	<b>3203</b>
Liveborn	3123
Other perinatal conditions	35
Hemolytic jaundice and perinatal jaundice	24
<b>3 Diseases of the circulatory system</b>	<b>2107</b>
Diseases of the heart	1443
Cerebrovascular disease	328
Hypertension	150
<b>4 Diseases of the digestive system</b>	<b>1295</b>
Lower gastrointestinal disorders	387
Biliary tract disease	190
Upper gastrointestinal disorders	159
<b>5 Diseases of the respiratory system</b>	<b>1145</b>
Respiratory infections	562
Asthma	164
Chronic obstructive pulmonary disease and bronchiectasis	154
<b>6 Injury and poisoning</b>	<b>978</b>
Complications	370
Fractures	310
Poisoning	58
<b>7 Neoplasms</b>	<b>855</b>
Benign neoplasms	237
Maintenance chemotherapy; radiotherapy	153
Secondary malignancies	77

In Greater Hobby Area, during the years 1999-2002, the most common causes of hospitalization were related to issues of childbirth and perinatal period conditions, cardiovascular and cerebrovascular diseases, digestive disorders, or respiratory infections/diseases.

Note that only the most common conditions are listed under each major category of diagnosis, and that the sum of these counts may not equal the total counts for the category.




<b>8 Diseases of the genitourinary system</b>	<b>692</b>
Diseases of the urinary system	383
Diseases of female genital organs	263
Diseases of male genital organs	46
<b>9 Symptoms; signs; and ill-defined conditions and factors influencing health status</b>	<b>644</b>
Factors influencing health care	397
Symptoms; signs; and ill-defined conditions	247
<b>10 Mental disorders</b>	<b>518</b>
Affective disorders	205
Schizophrenia and related disorders	124
Other psychoses	59

Data Source: Texas Department of State Health Services, Texas Health Care Information Collection

# Crime


The crime rate in urban areas is of concern to the residents, law enforcement and the local government. Crimes place stress on the residents of neighborhoods and affect their well-being. Of particular concern are violent crimes that threaten residents' lives, such as those involving firearms.

## Overview of Crime, 1999-2003



## Violent Crime, 1999-2003

The annual average rate of violent crime in Greater Hobby Area was 11.5 per 1,000 population, comparable to the Houston rate as a whole. The firearm-related violent crime rate in Greater Hobby Area was 4.5 per 1,000 population, slightly higher than the rate in Houston overall.



## Rate of Violent Crime by Super Neighborhood, 1999-2003

Greater Hobby Area was among the neighborhoods with the lowest annual average rates of violent crime in the city.

- Less than/Similar to the Houston rate
- Up to 25% above the Houston rate
- Up to 50% above the Houston rate
- Up to 75% above the Houston rate
- Greater than 75% above the Houston rate

Data Source: Houston Police Department

## Tuberculosis

Tuberculosis (TB) is caused by a specific type of bacteria that spreads from person to person through the air. TB typically affects the lungs but can also affect the brain and other organs. If this disease is left untreated it can be fatal.

From 1999 to 2003, 22 newly-acquired cases of tuberculosis were identified among residents of Greater Hobby Area, representing almost 2% of all cases diagnosed in Houston in that period. The annual average rate in Greater Hobby Area was 10.7 per 100,000 population, compared to 13.6 per 100,000 population in Houston as a whole. Both rates appeared much higher than the national Healthy People 2010 target of 1 case per 100,000 population.

The majority (77%) of these cases were among adults between 20 to 64 years of age.

Data Source: HDHHS, Bureau of TB Control

## Drowning and Submersion

Drowning and submersion injuries are often unintentional and are preventable through increased awareness of precautions that can be taken in and around bodies of water.

Fewer than 5 drowning or submersion cases were reported among Greater Hobby Area residents from 1999-2003.

Data Source: HDHHS, Bureau of Epidemiology

## Food-borne Diseases

Many food-related diseases are easily preventable. Eating well-cooked foods, keeping cooking areas free of contamination by thoroughly cleaning surfaces touched by raw meats and poultry, hand washing before handling food, and avoiding unpasteurized products are some of the measures that people can take to lower their risk of food-related disease.


Food-related diseases are typically under-reported. It is likely that many more cases occurred from 1999 to 2003 than were actually reported to health officials.

Typically Reported Diseases	Number of Cases
Hepatitis A	<5
Shigellosis	15
Salmonellosis	11
Campylobacteriosis	<5
Amebiasis	<5
Vibrio	<5

Data Source: HDHHS, Bureau of Epidemiology

# Environmental Health and Safety

Chemical emissions and waste released into the air, soil, and water can affect everyone. Knowing the locations and types of potential polluters allows residents to better monitor the potential environmental impact on their communities.



- Toxic Release Inventory (TRI) Facility
- ⊗ Major Storm Water Runoff Facility
- ◆ Hazardous Waste Treatment, Storage, or Disposal (TSD) Facility
- Large Quantity Generator (LQG) of Hazardous Waste
- Major Discharger of Air Pollutants
- ◆ Radioactive Waste Site
- ◆ Current Superfund Site
- ◆ Former Superfund Site
- ⊕ Active Landfill
- ⊕ Inactive Landfill
- ⊕ Closed Landfill
- Highway
- Major Roadway
- Railroad
- Bayou

## Regulated Facilities

The Environmental Protection Agency (EPA) and the Texas Commission on Environmental Quality (TCEQ) administer programs which monitor and regulate facilities with the potential to release significant amounts of hazardous chemicals to the environment.

Within one mile of Greater Hobby Area, there are 21 Toxic Release Inventory (TRI) reporting facilities, 9 Large Quantity Generators (LQG) of hazardous waste, 1 major discharger of air pollutants, 2 major storm water discharging facilities, 1 radioactive waste site, 1 former Superfund site on the TCEQ registry, and 1 current Superfund site on the EPA National Priorities List (NPL).

These facilities are regulated under one or more of the following federal statutes: the Emergency Planning and Community Right-to-Know Act (EPCRA), the Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA), the Resource Conservation and Recovery Act (RCRA), the Clean Air Act, and the Clean Water Act.



The EPA provides reports concerning federally regulated facilities through an online application called Envirofacts ([www.epa.gov/enviro/index.html](http://www.epa.gov/enviro/index.html)).

Type of Regulated Facility	Houston Count	Type of Regulated Facility	Houston Count
Toxic Release Inventory (TRI) Facilities (all reporting years)	302	Major Dischargers of Air Pollutants	71
Major Storm Water Runoff Facilities	56	Radioactive Waste Sites	4
Hazardous Waste Treatment, Storage, or Disposal (TSD) Facilities	35	Current Superfund Sites	12
Large Quantity Generators (LQG) of Hazardous Waste	132	Former Superfund Sites	5
		Active Landfills	9
		Inactive Landfills	2
		Closed Landfills	18

Data Sources: Environmental Protection Agency; Texas Commission on Environmental Quality

# HIV/AIDS

HIV (Human Immunodeficiency virus) attacks the immune system and can progress to Acquired Immune Deficiency Syndrome (AIDS). HIV is primarily transmitted through unprotected sex or sharing needles with someone infected with the virus. It can also be transmitted before or during birth and from breast milk from mother to child. Many of those infected are unaware of their HIV status, and therefore can transmit the disease unknowingly.




## New HIV Diagnosis Rate, 1999-2003

The annual average rate of new HIV diagnosis in the combined zip codes 77061 and 77075 (which include Greater Hobby Area) was 58% lower than the Houston-wide rate during the period 1999-2003; the rate of 26.7 cases per 100,000 population was far above the Healthy People 2010 goal of less than 1 new case per 100,000 population.


## HIV Risk Factors, 1999-2003

Sixty-nine percent of new HIV infections occurred in males in Greater Hobby Area. In one-quarter of all cases, the mode of transmission was unknown. Male-to-male sex accounted for about 33% of all reported cases. This was followed by heterosexual contact (25%) and use of IV drugs (14%). Three percent of new infections occurred in those reporting male-to-male sex and IV drug use.



## Rates of New HIV Diagnosis by Zip Code\*, 1999-2003

The annual average rates of new HIV diagnosis in zip codes 77061 and 77075 were each lower than that of most other zip codes in the city.



\* Annual average rates are calculated only for those zip codes that lie predominantly within the boundaries of the city of Houston.


Data Source: HDHHS, Bureau of Epidemiology

# Gonorrhea

Gonorrhea is a sexually transmitted disease (STD) caused by bacteria. If untreated, it can cause serious and permanent health problems in both women and men. It also places infected persons at greater risk for HIV. Though rare, it can result in death if untreated.



## New Gonorrhea Infection by Age, Sex, Race/Ethnicity

The annual average rate of new gonorrhea cases in Greater Hobby Area was slightly higher than the rate in Houston overall; both rates were much greater than the Healthy People 2010 goal of less than 19 cases per 100,000 population.




## Gonorrhea Infection by Age, Sex, Race/Ethnicity

Blacks, who represent 25% of Greater Hobby Area's population, accounted for 73% of new cases. Slightly more than half (54%) of all cases occurred in males, and persons aged 20-29 years accounted for most of the cases.



## Rates of Gonorrhea Infection by Super Neighborhood, 1999-2003


Greater Hobby Area was among those neighborhoods with elevated annual average rates of infection.



Data Source: HDHHS, Bureau of Epidemiology


# Syphilis

Syphilis is a sexually transmitted disease (STD) and is passed from person to person through direct contact with a syphilis sore. Sores occur mainly on the external genitals, vagina, anus, or in the rectum. Transmission occurs due to unprotected sex. The sores may also occur in lips and mouth. Untreated syphilis can progress into more serious conditions affecting the nervous system, heart and other organs, seriously impairing health.




**Rates of New Syphilis Infection, 1999-2003**

The annual average rate of new syphilis infection in Greater Hobby Area was slightly lower than the rate in Houston overall; both were far higher than the Healthy People 2010.



**Syphilis Cases by Age, Sex, Race/Ethnicity**

Nearly half (47%) of new cases in Greater Hobby Area occurred among Blacks. Slightly more females (53%) than males (47%) were affected by syphilis, and persons aged 30-49 years were the most affected group.



**Rates of Syphilis by Super Neighborhood, 1999-2003**

Greater Hobby Area was among the neighborhoods with the lowest annual average rates of infection.

- Less than/Similar to the Houston rate
- Up to 25% above the Houston rate
- Up to 50% above the Houston rate
- Up to 75% above the Houston rate
- Greater than 75% above the Houston rate
- Rate Unreliable


Data Source: HDHHS Bureau of Epidemiology

# Chlamydia

Chlamydia is the most frequently reported sexually transmitted disease (STD) in the nation. Women are more commonly screened for the infection than are men, and those 15 to 24 years of age appear to be the most affected, nation-wide. The symptoms are usually mild and not easily recognized, causing many with the infection not to seek treatment. If untreated, chlamydia can cause infertility in women.



## Rates of Chlamydia, 1999-2003

The annual average rate of chlamydia infection in Greater Hobby Area was 331 per 100,000 population, 16% higher than the rate in Houston overall. Although Black women between the ages of 15 and 24 years had the highest rate of infection in Greater Hobby Area, this rate was 26% lower than that of the same group in Houston overall.




## Chlamydia Infection By Age, Sex, and Race/Ethnicity, 1999-2003

Half of all cases in Greater Hobby Area occurred among Hispanics. Eighty-seven percent of all cases were female. Persons aged 20-29 years accounted for more than half (54%) of all cases.



## Rates of Chlamydia by Super Neighborhood, 1999-2003

Greater Hobby Area was among the neighborhoods with elevated annual average rates of infection.



Data Source: HDHHS Bureau of Epidemiology

## Technical Notes

The Community Health Profiles Project attempts to provide the most recent statistical information available on the health of communities. The 1999-2003 series represents a “baseline” against which changes in the health indicators of communities can be evaluated over time. Data used to compile this profile are derived from a variety of sources — local, state, and national. These data sources may collect information on different cycles and therefore gaps in available years of data may be observed within a single profile.

Except where noted otherwise, rates are calculated using 2000 census data for each community, including age, race, and sex distributions. Agreement between race/ethnicity classifications in the data used in this report and those derived from the census is imperfect; disease registries do not uniformly capture ethnicity along with race and categories of “Black”, “White,” “Asian,” and “Other” may overlap with “Hispanic” ethnicity. Despite potential overlap, in this profile, “Black” is meant as “non-Hispanic Black,” “White” as “non-Hispanic White,” and “Hispanic” as being persons of any race and of Hispanic/Latino culture and origin. The profiles group a range of years of data and present them, where most appropriate, as annual average incidence of the indicator. If the total number of events is less than five, the associated rate is considered unreliable and is not reported; however for Leading Causes of Death, the minimum number of deaths for reporting age-adjusted rates is set at 25. Statistics presented in profiles of super neighborhoods, medically-underserved areas (MUAs), and other geographies are based upon successful geocoding of the residence of individual cases within the boundaries of those geographic entities. The denominator in all cases is the year 2000 census, as the estimated “average” population for each year of the analysis period. Background Houston rates and Healthy People 2010 goals have been used for most indicators as a standard for comparison.

**Mortality data:** Mortality data have been obtained at the address level from the Texas Department of State Health Services for 1999-2003. The YPLL statistics are computed using 65 years of age as the end point. **Crime data:** Data for 1999-2003 have been acquired from the Houston Police Department at the address level of the site of the incident. **HIV/AIDS data:** As of this report, data were only available at the zip code level.

### Other notes

Data for a number of additional indicators considered important for a community’s assessment of its health and health planning efforts were not available at the time of printing of this document. These indicators, including various injury indicators, and more community-specific behavioral data are being collected or researched for potential inclusion in the future published version of this report.

## Community Health Profiles

Community-specific public health profiles on medically-underserved areas and the 88 super neighborhoods of Houston are available from the Houston Department of Health and Human Services at [www.houstontx.gov/health](http://www.houstontx.gov/health). Reports can also be requested by e-mail at [webadmin@cityofhouston.net](mailto:webadmin@cityofhouston.net), or by writing to:

### Community Health Statistics

Office of Surveillance & Public Health Preparedness  
Houston Department of Health and Human Services  
8000 N. Stadium Dr., 4th floor  
Houston, Texas 77054



City of Houston  
Department of Health and Human Services

Stephen L. Williams, M.Ed., M.P.A.  
Director

### Community Health Profiles

Produced by  
Community Health Statistics  
Office of Surveillance and Public Health  
Preparedness

Raouf Arafat, M.D., M.P.H.  
Assistant Director

Mark Perry, M.P.H.  
Editor

### **About Community Health Statistics (CHS)**

Community Health Statistics (CHS) is a program within the division of the Office of Surveillance and Public Health Preparedness of the Houston Department of Health and Human Services (HDHHS). It is comprised of epidemiologists, statisticians, and GIS analysts who acquire data through collaboration with multiple partners within and outside the department for analysis, interpretation, and sharing of information on local health issues.

Our mission is to serve the needs of HDHHS, and the needs of the scientific community, and general public as a resource for data and information on the indicators and the determinants of the health and well-being of geographically-defined communities, as well as of other distinct population groups within the city of Houston, Texas.